

Flows that are sums of hamiltonian cycles in abelian Cayley graphs

Joy Morris

University of Lethbridge
morris@cs.uleth.ca

David Petrie Moulton

Center for Communications Research
moulton@idaccr.org

Dave Witte*

Oklahoma State University
dwitte@math.okstate.edu
<http://www.math.okstate.edu/~dwitte>

Abstract


If X is any connected Cayley graph on any finite abelian group, we determine precisely which flows on X can be written as a sum of hamiltonian cycles. In particular, if the degree of X is at least 5, and X has an even number of vertices, then it is precisely the even flows, that is, the flows f , such that $\sum_{\alpha \in E(X)} f(\alpha)$ is divisible by 2. On the other hand, there are infinitely many examples of degree 4 in which not all even flows can be written as a sum of hamiltonian cycles. Analogous results were already known 10 years ago, from work of Brian Alspach, Stephen Locke, and Dave Witte, for the case where X is cubic, or has an odd number of vertices.

References

- B. Alspach, S. C. Locke, and D. Witte:
The Hamilton spaces of Cayley graphs on abelian groups,
Discrete Math. 82 (1990) 113–126.
- S. C. Locke and D. Witte:
Flows in circulant graphs of odd order are sums of Hamilton cycles,
Discrete Math. 78 (1989) 105–114.

Circulant graph. $\text{Circ}(12; \{3, 4\})$:

- $V(X) = \mathbb{Z}_{12}$
- edge $x \text{ --- } x \pm s$ for $s \in \{3, 4\}$.



Rem. $\text{Circ}(12; \{3, 4\}) \cong C_4 \square C_3$.

Abelian Cayley grf. Replace \mathbb{Z}_{12} with abel grp G .

Eg. $\text{Cay}(\mathbb{Z}_4 \oplus \mathbb{Z}_3; \{(1, 0), (0, 1)\})$
 $\cong C_4 \square C_3$.

Always assume Cayley graphs are connected.

I.e., S generates G .

Recall. Any flow in any graph is a sum of cycles.

Thm (Alspach, Locke, Witte, 1990).

Every flow in any abelian Cayley graph X
= sum of hamiltonian cycles
(unless $X \cong C_{\text{odd}} \square K_2$)
(mod 2).

Thm (Locke, Witte, 1989).

Every flow in any abelian Cayley graph X
= sum of hamiltonian cycles
if X has odd order
(unless $X \cong K_3 \square K_3$).

Locke-Witte also settled the cubic graphs.

Remains: graphs of even order, with degree ≥ 4 .

X has even order \Rightarrow every ham cyc is **even** flow,

i.e., $\sum_{e \in E(X)} f(e)$ is even.


So sum of ham cycs is always an even flow.

Thm (Morris, Moulton, Witte).
 Even flow in any abel Cayley grf X of even order
 = sum of hamiltonian cycles
 if $\deg X \geq 5$.

Thm (M, M, W). $X =$ abel Cay grf of deg 4.
 Every even flow is sum of ham cycles unless

- $4 \nmid |V(X)|$ and X is not bipartite; or
- $X =$ square of cycle.


Defn. Square of cycle = $\text{Cay}(\mathbb{Z}_n; \{1, 2\})$




Rem. We are still working on the converse.

- $\mathcal{E} = \{\text{even flows on } X\}$
- $\mathcal{H} = \{\text{sums of hamiltonian cycles}\}$


Basic 4-cycle. $s, t \in S \Rightarrow (s, t, s^{-1}, t^{-1}) \in \mathcal{F}$




Lem. If every basic 4-cycle is in \mathcal{H} ,
 then $\mathcal{E} \subset \mathcal{H}$.




Cor. If $\mathcal{H} \ni$ sum of odd number of basic 4-cycles,
 then every basic 4-cycle is in \mathcal{H} ,
 so $\mathcal{E} \subset \mathcal{H}$.



Prop. Some ham cyc in $P_3 \square P_3 \square P_2$
 is the sum of an odd number of 4-cycles



Prop. Some ham cyc in $P_3 \square P_3 \square P_2$
 is the sum of an even number of 4-cycles



Cor. If $P_3 \square P_3 \square P_2 \subset X$,
 then $\mathcal{E} \subset \mathcal{H}$.

Cor. If $\deg X \geq 5$, then $\mathcal{E} \subset \mathcal{H}$.